



Contents lists available at ScienceDirect

# Journal of Science and Medicine in Sport

journal homepage: [www.elsevier.com/locate/jsams](http://www.elsevier.com/locate/jsams)



Original research

## Incidence and predictors of exertional hyperthermia after a 15-km road race in cool environmental conditions

Matthijs T.W. Veltmeijer<sup>a,1</sup>, Thijs M.H. Eijsvogels<sup>a,b,1</sup>, Dick H.J. Thijssen<sup>a,c</sup>,  
Maria T.E. Hopman<sup>a,\*</sup>

<sup>a</sup> Department of Physiology, Radboud University Medical Centre, the Netherlands

<sup>b</sup> Henry Low Heart Centre, Department of Cardiology, Hartford Hospital, USA

<sup>c</sup> Research Institute for Sport and Exercise Science, Liverpool John Moores University, United Kingdom

### ARTICLE INFO

#### Article history:

Received 23 December 2013  
Received in revised form 10 April 2014  
Accepted 17 April 2014  
Available online xxx

#### Keywords:

Heatstroke  
Heat stress disorders  
Body temperature regulation  
Exercise  
Body temperature

### ABSTRACT

**Objectives:** Current knowledge about the incidence and risk factors for exertional hyperthermia (core body temperature  $\geq 40^\circ\text{C}$ ) is predominantly based on military populations or small-sized studies in athletes. We assessed the incidence of exertional hyperthermia in 227 participants of a 15-km running race, and identified predictors for exertional hyperthermia.

**Design:** Observational study.

**Methods:** We measured intestinal core body temperature before and immediately after the race. To identify predictive factors of maximum core body temperature, we entered sex, age, BMI, post-finish dehydration, number of training weeks, fluid intake before and during the race, finish time, and core body temperature change during warming-up into a backward linear regression analysis. Additionally, two subgroups of hyperthermic and non-hyperthermic participants were compared.

**Results:** In a WBGT of  $11^\circ\text{C}$ , core body temperature increased from  $37.6 \pm 0.4^\circ\text{C}$  at baseline to  $37.8 \pm 0.4^\circ\text{C}$  after warming-up, and  $39.2 \pm 0.7^\circ\text{C}$  at the finish. A total of 15% of all participants had exertional hyperthermia at the finish. Age, BMI, fluid intake before the race and the core body temperature change during warming-up significantly predicted maximal core body temperature ( $p < 0.001$ ). Participants with hyperthermia at the finish line had a significantly greater core body temperature rise ( $p < 0.01$ ) during the warming-up compared to non-hyperthermic peers, but similar race times ( $p = 0.46$ ).

**Conclusion:** 15% of the recreational runners developed exertional hyperthermia, whilst core body temperature change during the warming-up was identified as strongest predictor for core body temperature at the finish. This study emphasizes that exertional hyperthermia is a common phenomenon in recreational athletes, and can be partially predicted.

© 2014 Sports Medicine Australia. Published by Elsevier Ltd. All rights reserved.

### 1. Introduction

Current knowledge about the incidence and risk factors for exertional hyperthermia (core body temperature (CBT)  $\geq 40^\circ\text{C}$ )<sup>1</sup> and heat illness is largely based on retrospective studies investigating military populations during military exercises.<sup>2–4</sup> These studies involved well-trained soldiers performing continuous exercise (e.g. long-distance running) superimposed by bouts of high-intensity anaerobic exercise (e.g. heavy lifting). This type of exercise is substantially different from a typical athletic event popular in the

general public, during which the athletes typically only perform continuous high-intensity exercise. Furthermore, paramount to the general public is that it is characterized by a wide range of individual traits, including a wide diversity in body mass, age and training and health status.<sup>5</sup> As all these factors may affect thermoregulatory responses differently, each individual may be subject to a different risk for developing heat-related problems, such as exertional hyperthermia or heat illness.<sup>6–8</sup> Previous studies that did focus on thermoregulation in participants of athletic events or outdoor time trials in cool to moderate conditions are based on relatively small to moderate sample sizes.<sup>9–11</sup> These studies reported exertional hyperthermia in 0–23% of their participants, mostly after performing marathon races, and this wide range makes it tenuous to draw any firm conclusions. In addition, no previous authors have confirmed whether this knowledge is applicable to the general public

\* Corresponding author.

E-mail address: [Maria.Hopman@Radboudumc.nl](mailto:Maria.Hopman@Radboudumc.nl) (M.T.E. Hopman).

<sup>1</sup> These authors contributed equally to the manuscript.

based on measurements in a large and heterogeneous sample size. Based on previous literature, risk factors for heat illness such as metabolic rate (i.e. running speed),<sup>9,12,13</sup> dehydration at the finish line and low fluid intake before and during the race,<sup>1,14,15</sup> increased body mass index,<sup>1,4,16</sup> poor training,<sup>4</sup> advanced age<sup>14</sup> and male sex<sup>17</sup> could significantly predict CBT at the finish line. However, it has never been confirmed whether these risk factors can be applied to identify athletes at risk of developing hyperthermia during large sports events.

Therefore, the main purpose of this study was to assess the CBT in a large ( $n=227$ ) heterogeneous group of participants of a 15-km running race (Seven Hills Run, Nijmegen, the Netherlands). This is one of the largest running events held in the Netherlands (>30,000 participants), and holds the men's and women's 15-km world records set in 2010 and 2009 respectively. Secondly, we identified factors that significantly predicted CBT at the finish line using a backward linear regression analysis. Thirdly, we aimed to assess the differences between athletes finishing with a high CBT versus those with a low CBT, in order to identify key features that may explain the CBT rise during exercise. To that end, we compared body and race characteristics in athletes with a finish CBT  $\geq 40^\circ\text{C}$  (hyperthermic athletes) to an equally sized group of athletes that finished with the lowest CBT of all participants (non-hyperthermic athletes).

## 2. Methods

Five-hundred participants of the Seven Hills Run were randomly contacted and, if interested, were sent a study protocol. All volunteers were screened for the presence of any exclusion criteria for using the temperature pill: (1) a history of obstructive or inflammatory bowel disease, or any prior abdominal surgery, (2) the presence of any implanted electric (medical) device, (3) a scheduled MRI

scan within 1 week after the event, or (4) pregnancy. Two hundred twenty-seven participants were included in the study: 111 men and 116 women were aged  $45 \pm 11$  years and had a BMI of  $22.7 \pm 2.7 \text{ kg/m}^2$  (Table 1). Study procedures were approved by the Radboud University Medical Centre Ethics Committee, accorded to the principles of the Declaration of Helsinki, and all participants provided written informed consent before participation.

Prior to the race, participants completed a questionnaire pertaining to their physical training. Participants self-reported their fluid intake from the time of getting out of bed on the day of the race and during the race. Body weight was measured before and after the race in a laboratory set up 50 m from the finish line. CBT was measured at baseline in the laboratory about 2 h before the start, 1 min before the start (i.e. after warming-up), and within 15 s after finishing. Due to the large total number of participants in the race, runners started the race phased into in 9 separate 'waves' over a 1 h period. Ten research assistants measured CBT in  $25 \pm 1$  participants per wave using 5 wireless receivers. Participants with a CBT  $\geq 40^\circ\text{C}$  upon finishing were compared to an equal number of participants that finished with the lowest CBT.

Participants ingested an individually calibrated telemetric temperature pill at least 5 h (8 a.m.) before the race (start 1 p.m.) to prevent interaction of the CBT measurements with fluid ingestion during testing.<sup>18</sup> CBT was measured using a portable telemetry system (CorTemp™ system, HQ Inc., Palmetto, USA). This measuring system has been demonstrated to safely and reliably measure CBT.<sup>19,20</sup> The average of three consecutive measurements for each time point was used for further analyses. The change in CBT during warming-up was calculated by subtracting CBT at baseline from the CBT before the start of the race.

Body weight was measured to the nearest 0.1 kg using an automatically calibrated balance (Seca 888; Hamburg, Germany) before and within 10 min after the race. The relative change in body

**Table 1**  
Participant demographics, physical activity pattern, race characteristics, CBT, body weight, and fluid intake in the total group, a subgroup of hyperthermic participants at the finish line and a subgroup of non-hyperthermic participants.

	Total group	Exertional hyperthermia	Non-hyperthermia	p-Value
<b>Characteristics</b>				
Sex (male: female)	111:116	15:16	16:15	0.80
Age (years)	$45 \pm 11$	$43 \pm 11$	$45 \pm 11$	0.49
Body mass index ( $\text{kg/m}^2$ )	$22.7 \pm 2.7$	$23.0 \pm 2.4$	$22.1 \pm 2.6$	0.17
<b>Physical activity pattern</b>				
Number of previous participations in this event	$5 \pm 4$	$4 \pm 3$	$5 \pm 4$	0.30
Training (weeks)	28 (range 0–52)	30 (range 3–52)	24 (range 0–52)	0.32
Running exercise (sessions/week)	$2.5 \pm 0.9$	$2.4 \pm 0.8$	$2.7 \pm 1.0$	0.21
<b>Race characteristics<sup>a</sup></b>				
Split time 0–5 km (min)	$26.2 \pm 3.7$	$25.8 \pm 3.0$	$25.4 \pm 4.0$	ANOVA (split times): Time/group/time $\times$ group $p = 0.16/p = 0.46/p = 0.10$
Split time 5–10 km (min)	$26.4 \pm 3.9$	$26.0 \pm 3.1$	$25.5 \pm 4.0$	
Split time 10–15 km (min)	$26.6 \pm 6.7$	$26.4 \pm 3.5$	$25.3 \pm 4.1$	
Total race time (min)	$79.1 \pm 12.8$	$78.3 \pm 9.4$	$76.2 \pm 11.9$	
Total race speed ( $\text{km/h}$ )	$11.7 \pm 1.8$	$11.7 \pm 1.5$	$12.1 \pm 1.9$	
<b>Core body temperature</b>				
Baseline	$37.6 \pm 0.4$	$37.6 \pm 0.4$	$37.4 \pm 0.4$	0.26
CBT change during warming-up	$0.2 \pm 0.5$	$0.5 \pm 0.5$	$0.1 \pm 0.4$	<0.01
Start	$37.8 \pm 0.4$	$38.0 \pm 0.5$	$37.5 \pm 0.3$	<0.001
Finish	$39.2 \pm 0.7$	$40.4 \pm 0.4$	$38.1 \pm 0.3$	–
<b>Body weight</b>				
Baseline body weight (kg)	$71.7 \pm 11.9$	$72.3 \pm 11.4$	$70.0 \pm 12.8$	0.46
Finish body weight (kg)	$70.8 \pm 11.7$	$71.3 \pm 11.6$	$69.0 \pm 12.6$	0.48
$\Delta$ Body weight (%)	$-1.5 \pm 0.6$	$-1.7 \pm 0.6$	$-1.5 \pm 0.6$	0.39
Classifying as dehydrated <sup>b</sup> (%)	21	29	24	0.67
<b>Fluid intake</b>				
Fluid intake before race (L)	$1.18 \pm 0.47$	$1.0 \pm 0.4$	$1.3 \pm 0.5$	<0.05
Fluid intake during race (L)	$0.06 \pm 0.12$	$0.04 \pm 0.06$	$0.07 \pm 0.13$	0.24

<sup>a</sup> Differences in split times were tested using a Two-Way Repeated Measures ANOVA.

<sup>b</sup> Participants classified as 'dehydrated' if body weight at the finish line was reduced  $\geq 2\%$ .

weight was calculated and dehydration was defined as a body weight loss of  $\geq 2\%$ .<sup>21</sup> Participants were allowed to drink ad libitum before and during the race, and self-reported the time and amount (standard sized cups, bottles, etc.) of their individual fluid intake before and during the race. No restrictions were imposed on the type of fluids consumed, though participants were requested to refrain from drinking between finishing and the second body weight measurement to avoid overestimating the post-race body weight. Furthermore, body weight change during the race (delta body weight) was corrected for fluid intake during the race by adding the fluid intake to the delta body weight.

Individual split times after 5-, 10- and 15-km were obtained from the organizational measuring system (ChampionChip®, MYLAPS, Nijmegen, the Netherlands).

Wet bulb globe temperature (WBGT) was measured every 30 min throughout the day using a portable climate monitoring device (Davis Instruments Inc., Hayward, USA) positioned in the start/finish area.

Statistical analyses were performed using the Statistical Package for the Social Sciences (IBM SPSS Statistics for Windows, Version 20.0. IBM Corp., Armonk, NY, USA). Data was reported as mean  $\pm$  standard deviation unless otherwise indicated. A backward linear regression analysis was used to identify factors that significantly predicted finish CBT. Age, sex, BMI, finish time, fluid intake before and during the race, the presence of post-finish dehydration, the delta CBT between baseline and the start and the number of training weeks were included as potential factors that could predict finish CBT. Differences between the subgroups of hyperthermic and non-hyperthermic participants were tested using a Student's *t*-test for the continuous data, and a chi-square test for the nominal data (i.e. the presence of post-finish dehydration). The significance level was set at  $p \leq 0.05$ .

### 3. Results

The average race time was  $79 \pm 13$  min (range 55–165 min), with a mean running speed of  $11.7 \pm 1.8$  km/h (Table 1). Furthermore, the split times after every 5-km point were comparable ( $p=0.33$ ) across the race. Under cool environmental conditions (WBGT was stable at  $11^\circ\text{C}$  throughout,  $T_{\text{DRY-BULB}}$   $10.5^\circ\text{C}$ , relative humidity 87%), CBT increased from  $37.6 \pm 0.4^\circ\text{C}$  at baseline to  $37.8 \pm 0.4^\circ\text{C}$  after the warming-up at the start, and was  $39.2 \pm 0.7^\circ\text{C}$  upon finishing (Fig. 1). CBT could not be measured at the finish line in 18 participants (8%). Thirty-one participants (15%) showed a CBT  $\geq 40^\circ\text{C}$  and were hence classified as being hyperthermic. None of the participants reported any apparent heat-related physical complaints.

Between baseline and finish, a  $1.5 \pm 0.6\%$  reduction in total body weight was observed, and 21% of all participants were classified as dehydrated ( $\geq 2\%$  decrease in body weight). Self-reported fluid




Fig. 1. Frequency distribution of core body temperature (CBT) at baseline (light bars) and finish line (dark bars). Fifteen percent of the participants had exertional hyperthermia at the finish line after completion of the race.

intake before the start of the race was  $1.18 \pm 0.47$  L, whilst intake was  $0.06 \pm 0.12$  L during the race.

The backward linear regression model ( $r=0.41$ ,  $p<0.001$ ) identified age ( $B=-0.01$ ,  $p=0.03$ ), BMI ( $B=0.06$ ,  $p<0.01$ ), self-reported fluid intake before the race ( $B=-0.30$ ,  $p=0.02$ ) and CBT change during warming-up ( $\beta=0.56$ ,  $p<0.001$ ) as parameters that significantly predicted CBT at the finish line (Table 2). Sex, finish time, self-reported fluid intake during the race, the presence of post-finish dehydration and number of training weeks were also entered into the regression analysis, but did not appear to influence finish CBT. These results remained unchanged when the analysis was repeated after replacing the total finish time for the split time in the last 5-km.

CBT change during warming-up was identified as the strongest predictor in our model. Accordingly, we were interested in the risk stratification of participants that demonstrated a CBT change during warming-up that was 2 times higher than the group average ( $0.2^\circ\text{C}$ ). Therefore, we created a new dichotomous variable in which participants were classified to a CBT rise  $\leq 0.4^\circ\text{C}$  or  $>0.4^\circ\text{C}$  during warming-up. Exertional hyperthermia occurred significantly ( $p<0.001$ ) more frequent in the participants with a CBT rise  $>0.4^\circ\text{C}$  (33.3%) compared to participants with a CBT rise  $\leq 0.4^\circ\text{C}$  (9%) during warming-up (OR: 5.1, 95% CI: 2.2–11.7).

Two subgroups comprising 31 hyperthermic participants and 31 participants that finished with the lowest CBT of the total group (non-hyperthermic group) were selected for additional analyses

Table 2  
 Predictors for finish CBT.

Variable	Correlation Coefficients			Linear Regression analysis <sup>a</sup>		
	B	95% CI	$\beta$	B	95% CI	$\beta$
Constant				38.7	37.7–39.7	
Age	-0.01	-0.02 to -0.001	-0.1 <sup>c</sup>	-0.01	-0.02 to -0.001	-0.1 <sup>c</sup>
BMI	0.07	-0.02 to 0.12	0.25 <sup>b</sup>	0.06	0.02–0.10	0.21 <sup>b</sup>
CBT rise after warming-up	0.54	0.30–0.78	0.33 <sup>a</sup>	0.56	0.33–0.80	0.35 <sup>a</sup>
Fluid intake before race	-0.28	-0.52 to -0.03	-0.1 <sup>c</sup>	-0.30	-0.53 to -0.06	-0.1 <sup>c</sup>

<sup>a</sup>  $R^2$  for model = 0.167; adjusted  $R^2$  = 0.147.

<sup>a</sup>  $p < 0.001$ .

<sup>b</sup>  $p < 0.01$ .

<sup>c</sup>  $p < 0.05$ .

NS, not significant.

CI = confidence interval;  $\beta$  = standardized B.

with respect to body and race characteristics (Table 1). Within these groups, no differences were found for age, BMI, physical activity, race times, or body weight changes. The CBT change between baseline and the start of the race was significantly greater in the hyperthermic participants compared to the non-hyperthermic group ( $p < 0.01$ ). Furthermore, self-reported fluid intake before the race was significantly higher in the non-hyperthermic participants compared to the hyperthermic participants ( $p < 0.05$ ), whilst self-reported fluid intake during the race was similar in both groups.

#### 4. Discussion

This study assessed the incidence of exertional hyperthermia in a large and heterogeneous group of athletes during a 15-km running event. We found that 15% of our participants passed the finish line with a CBT  $\geq 40^\circ\text{C}$ . Taking into consideration that approximately 30,000 participants entered the race, this would mean that as many as 4200 participants may have developed exertional hyperthermia. We found that age, BMI, CBT change during warming-up and self-reported fluid intake before the race predicted CBT at the finish line, whereas sex, self-reported fluid intake during the race, the presence of post-finish dehydration, and the number of training weeks had no impact on finish CBT. Interestingly, we found no differences in body and race characteristics between the subgroups of hyperthermic athletes and their non-hyperthermic peers, apart from a higher (0.3 L) fluid intake before the race by the non-hyperthermic athletes. However, the CBT change after the warming-up was significantly greater, and self-reported fluid intake before the race was significantly lower in the hyperthermic participants compared to the non-hyperthermic participants. Additionally, we found that athletes with a CBT rise after warming-up  $>0.4^\circ\text{C}$  were significantly more likely (OR 5.1) to develop hyperthermia at the finish line. These results suggest that exertional hyperthermia is a common phenomenon in recreational athletes, does not necessarily result in physical complaints or a reduced exercise performance, and can be partially predicted.

To the best of our knowledge, this is the first study to measure CBT at the finish line in a large and heterogeneous group of runners. We found that 15% of our participants developed a CBT  $\geq 40^\circ\text{C}$  after 15-km running in cool conditions. Previous smaller-sized studies demonstrated that exertional hyperthermia occurs frequently during military exercises and athletic events,<sup>10,16,22</sup> and our results are the first that confirm the relatively high incidence in a large sample-size from the general public. Interestingly, the reported incidence of hyperthermia in the present (15%) and previous studies (50% and 56%)<sup>10,22</sup> seems to be higher than the reported CBT in several other studies (0%, 3% and 11%).<sup>11,12,23</sup> The difference may relate to cooler conditions or longer exercise duration that was possibly performed at lower exercise intensities in the studies that found a smaller incidence. Another potential explanation might relate to the use of rectal probes in the three latter studies, which were inevitably inserted several minutes after finishing the race. As passive cooling may result in a CBT drop of 0.2–0.5  $^\circ\text{C}$  within the first 5 min post-exercise,<sup>24</sup> the studies measuring rectal temperatures may have in fact underestimated the actual CBT in their participants.

Despite the high incidence of exertional hyperthermia, none of our participants reported any apparent heat-related complaints. Furthermore, we found no differences in race times between the subgroups with and without exertional hyperthermia, nor any differences in split times at the 5-, 10- and 15-km points (i.e. a stable running pace throughout the race). These results suggest that hyperthermia per se does not necessarily result in a reduced exercise performance. Previous authors have questioned the presence of a 'critical' CBT threshold for a reduced exercise performance,<sup>10,25</sup>

the latter of which is thought to be caused by a neurologically-mediated sustained decrease in muscle force production.<sup>26</sup> Based on a substantially larger sample-size, our results raise the same hesitations. Tolerance for hyperthermia is widely variable amongst athletes, but was demonstrated to be better in ambient conditions that favour low skin temperatures.<sup>10,25,27</sup> The relatively cool environmental conditions in the present study may therefore at least partially explain why both the hyperthermic and non-hyperthermic participants were able to preserve their race times throughout the race. It would be of great interest to further explore this by assessing CBT and skin temperature in a large group of athletes during a similar running event under cool, moderate and hot environmental conditions, and investigate whether hyperthermia only leads to reduced performance levels when skin temperatures are high.

Interestingly, participants who showed a CBT rise  $>0.4^\circ\text{C}$  between baseline and the start of the race (i.e. after warming-up) had a significantly higher occurrence (OR 5.1) of exertional hyperthermia upon finishing than participants who had a lower CBT rise during warming-up. Furthermore, the hyperthermic participants had a significantly greater CBT rise during the warming-up compared to their non-hyperthermic peers. This finding raises the question as to whether a strong CBT rise during warming-up can help in identifying athletes at risk of developing hyperthermia during the race. To the best of our knowledge, no such relationship has been reported previously. However, it has been reported that a previous episode of heat illness predisposes athletes for a repeat event,<sup>1</sup> suggesting that an intrinsic predisposition for heat illness may exist. This hypothesis is reinforced by our finding that athletes with the strongest CBT rise after the warming-up were significantly more likely to become hyperthermic at the finish line, which suggests that a similar intrinsic predisposition for developing hyperthermia may exist as well. The clinical implication of this finding is that measuring the CBT rise during warming-up may aid in identifying individuals who should be monitored more carefully, and may help initiate a more direct treatment strategy if problems do occur (i.e. physical complaints or performance detriments). Furthermore, it might also help to identify individuals who are most likely to benefit from pre-cooling interventions.

By measuring a large number of possible predictive factors for the finish CBT, we were able to predict 16.7% of the total finish CBT. The finding that a higher self-reported fluid intake before the race was predictive for a lower finish CBT in the total group (regression analysis) and the non-hyperthermic subgroup versus the hyperthermic subgroup could relate to a difference in hydration status prior to the exercise bout. Since we did not measure the participants' hydration status before the start of the race (e.g. through measuring urinary specific gravity at baseline), it is difficult to draw any definitive conclusions about this. However, previous literature recommended that the consumption of ~0.4–0.5 L of fluids 4 h before the start of exercise ensures euhydration at the start of exercise.<sup>28</sup> Since the participants of the present study consumed  $1.2 \pm 0.5$  L of fluids in the ~6 h time span prior to the race, it is likely that they started the race in a euhydrated state. Apart from the potential role of fluid intake before the race, we found no other differences between both subgroups of hyperthermic and non-hyperthermic participants. This emphasizes the difficulty to predict CBT during exercise using traditional measures. Previous studies have shown that exercise intensity and physical fitness affect CBT during exercise.<sup>1,4,13</sup> As our study was performed under race conditions, most participants performed in the upper range of their possible exercise intensity levels. This leaves only a small range of exercise intensities to correlate with CBT, and may explain why we found no significant relationship between both parameters.

The strengths of this study are the inclusion of a large and heterogeneous group of participants, the real-life race setting in which

participants reach peak performances that cannot be simulated in a laboratory situation, and the fact that we measured CBT within seconds after finishing. However, this study was limited by the fact that no correction was applied for sweat entrapment in the participants' clothing after the race concerning the body weight measurement after the race. Based on previous literature, it is known that not correcting for sweat entrapment in clothing may lead to a measurement error of 10%.<sup>9</sup> As subjects in the present study had an average weight loss of  $-1.0 \pm 0.5$  kg, actual sweat losses might have been underestimated by  $\sim 0.10$  kg on average. However, since the average change in body weight in the present study was already limited and remained below the recommended maximal weight change of 2%,<sup>21</sup> we believe that this small underestimation did not substantially influence our conclusions.

## 5. Conclusion

In cool environmental conditions (WBGT 11 °C), 15% of the participants passed the finish line with exertional hyperthermia. The stable and similar race times in both the hyperthermic and non-hyperthermic participants suggest that hyperthermia per se does not necessarily result in performance detriments. The limited value of the predictors of the increase in CBT and the great similarities between both subgroups of hyperthermic and non-hyperthermic participants, emphasizes the difficulty to predict CBT during exercise. However, we did find that age, BMI, self-reported fluid intake before the race and CBT rise during warming-up significantly predicted finish CBT, and that participants with a CBT rise after the warming-up  $>0.4$  °C were significantly more likely to develop hyperthermia at the finish line (OR 5.1), which may provide new prospects for predicting CBT and associated heat related problems during exercise.

## Practical implications

- Athletes who show sharp rises in their body temperature at an early stage of exercise may be at increased risk of developing exertional hyperthermia. This finding could aid in identifying any athletes who are at increased risk, so that they can be monitored more carefully if desired.
- Furthermore, identifying athletes with the sharpest rises in body temperature during warming-up may also aid in identifying those athletes that may benefit the most from cooling interventions.
- We provide evidence that hyperthermia does not necessarily result in a reduced exercise performance or lead to health problems. This implies that no immediate action needs to be taken when an athlete is hyperthermic without presenting with physical complaints or performance detriments.

## Acknowledgements

We thank the following persons for their invaluable assistance with the collection of data: S. van Dartel, R. Meurders, M. Sander, R. Scholten, P. de Groot, N. van Duijnhoven, T. Schreuder, B. Hijmans-Kersten, J. Evers and 40 other volunteers during this study.

Furthermore, we thank the organization of the 'Seven Hills Run' for providing facilities to set up our on-site laboratory.

This study was funded by the Department of Physiology of the Radboud University Medical Centre.

Dr. Eijsvogels received a Rubicon Grant (825.12.016). Dr. Thijsen is financially supported by the Netherlands Heart Foundation (E. Dekker-stipend 2009T064).

## References

1. Armstrong LE, Casa DJ, Millard-Stafford M et al. American College of Sports Medicine position stand Exertional heat illness during training and competition. *Med Sci Sports Exerc* 2007; 39(3):556–572.
2. Carter 3rd R, Chevront SN, Williams JO et al. Epidemiology of hospitalizations and deaths from heat illness in soldiers. *Med Sci Sports Exerc* 2005; 37(8):1338–1344.
3. Rav-Acha M, Hadad E, Epstein Y et al. Fatal exertional heat stroke: a case series. *Am J Med Sci* 2004; 328(2):84–87.
4. Wallace RF, Kriebel D, Punnett L et al. The effects of continuous hot weather training on risk of exertional heat illness. *Med Sci Sports Exerc* 2005; 37(1):84–90.
5. Taylor BA, Zaleski AL, Capizzi JA et al. Influence of chronic exercise on carotid atherosclerosis in marathon runners. *BMJ Open* 2014; 4(2):e004498.
6. Bouchama A, Knochel JP. Heat stroke. *N Engl J Med* 2002; 346(25):1978–1988.
7. Coris EE, Ramirez AM, Van Durme DJ. Heat illness in athletes: the dangerous combination of heat, humidity and exercise. *Sports Med* 2004; 34(1):9–16.
8. Simon HB. Hyperthermia. *N Engl J Med* 1993; 329(7):483–487.
9. Chevront SN, Haymes EM. Thermoregulation and marathon running: biological and environmental influences. *Sports Med* 2001; 31(10):743–762.
10. Ely BR, Ely MR, Chevront SN et al. Evidence against a 40 degrees C core temperature threshold for fatigue in humans. *J Appl Physiol* (1985) 2009; 107(5):1519–1525.
11. Maughan RJ. Thermoregulation in marathon competition at low ambient temperature. *Int J Sports Med* 1985; 6(1):15–19.
12. Maughan RJ, Leiper JB, Thompson J. Rectal temperature after marathon running. *Br J Sports Med* 1985; 19(4):192–195.
13. Noakes TD, Myburgh KH, du Plessis J et al. Metabolic rate, not percent dehydration, predicts rectal temperature in marathon runners. *Med Sci Sports Exerc* 1991; 23(4):443–449.
14. Casa DJ, Armstrong LE, Kenny GP et al. Exertional heat stroke: new concepts regarding cause and care. *Curr Sports Med Rep* 2012; 11(3):115–123.
15. Sawka MN. Physiological consequences of hypohydration: exercise performance and thermoregulation. *Med Sci Sports Exerc* 1992; 24(6):657–670.
16. Bedno SA, Li Y, Han W et al. Exertional heat illness among overweight U.S Army recruits in basic training. *Aviat Space Environ Med* 2010; 81(2):107–111.
17. Howe AS, Boden BP. Heat-related illness in athletes. *Am J Sports Med* 2007; 35(8):1384–1395.
18. Wilkinson DM, Carter JM, Richmond VL et al. The effect of cool water ingestion on gastrointestinal pill temperature. *Med Sci Sports Exerc* 2008; 40(3):523–528.
19. Byrne C, Lim CL. The ingestible telemetric body core temperature sensor: a review of validity and exercise applications. *Br J Sports Med* 2007; 41(3):126–133.
20. Gant N, Atkinson G, Williams C. The validity and reliability of intestinal temperature during intermittent running. *Med Sci Sports Exerc* 2006; 38(11):1926–1931.
21. Casa DJ, Clarkson PM, Roberts WO. American College of Sports Medicine roundtable on hydration and physical activity: consensus statements. *Curr Sports Med Rep* 2005; 4(3):115–127.
22. Byrne C, Lee JK, Chew SA et al. Continuous thermoregulatory responses to mass-participation distance running in heat. *Med Sci Sports Exerc* 2006; 38(5):803–810.
23. Pugh LG, Corbett JL, Johnson RH. Rectal temperatures, weight losses, and sweat rates in marathon running. *J Appl Physiol* 1967; 23(3):347–352.
24. Casa DJ, Armstrong LE, Ganio MS et al. Exertional heat stroke in competitive athletes. *Curr Sports Med Rep* 2005; 4(6):309–317.
25. Chevront SN, Kenefick RW, Montain SJ et al. Mechanisms of aerobic performance impairment with heat stress and dehydration. *J Appl Physiol* (1985) 2010; 109(6):1989–1995.
26. Nielsen B, Nybo L. Cerebral changes during exercise in the heat. *Sports Med* 2003; 33(1):1–11.
27. Schlader ZJ, Simmons SE, Stannard SR et al. Skin temperature as a thermal controller of exercise intensity. *Eur J Appl Physiol* 2011; 111(8):1631–1639.
28. American College of Sports Medicine, Sawka MN, Burke LM et al. American College of Sports Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc* 2007; 39(2):377–390.